

# A Level Pure Mathematics

## Practice Test 3: Exponentials and Logarithms

### Instructions:

Answer all questions. Show your working clearly.

Calculators may be used unless stated otherwise.

Time allowed: 2 hours

### Section A: Laws of Indices and Exponential Functions

1. Simplify these expressions using laws of indices:

(a)  $5^2 \times 5^6 \times 5^{-4}$

(b)  $\frac{7^8 \times 7^{-4}}{7^3}$

(c)  $(4^2)^5 \times 4^{-6}$

(d)  $\frac{(5^2)^3 \times 5^{-3}}{5^2}$

(e)  $(4^3 \times 4^{-1})^2$

(f)  $\frac{16^{2x} \times 16^{x-3}}{16^{3x+1}}$

2. Solve these exponential equations:

(a)  $5^x = 125$

(b)  $4^{x+2} = 64$

(c)  $2^{3x-1} = 32$

(d)  $8^x = \frac{1}{4}$

(e)  $25^x = 5^{x+6}$

(f)  $4^{2x} = 8^{x-3}$

3. Express in the form  $a^x$  where  $a$  is a rational number:

(a)  $(\frac{1}{5})^x \times 25^x$

(b)  $\frac{32^x}{8^{2x}}$

(c)  $(125)^{\frac{x}{3}} \times (\frac{1}{5})^x$

(d)  $\frac{49^x \times 7^{-3x}}{343^{\frac{x}{3}}}$

4. Sketch the graphs of these exponential functions, showing key features:

(a)  $y = 5^x$

(b)  $y = (\frac{1}{5})^x$

(c)  $y = 4^x + 2$

(d)  $y = 5^{x-2}$

(e)  $y = -5^x$

(f)  $y = 5^{-x}$

5. For the function  $f(x) = 3e^x$ :

(a) State the domain and range

(b) Find the y-intercept

(c) Describe the behavior as  $x \rightarrow \infty$  and  $x \rightarrow -\infty$ (d) Find  $f'(x)$  and comment on the gradient(e) Sketch the graph, showing the tangent at  $(0, 3)$ 

## Section B: Logarithmic Functions and Properties

6. Express these in logarithmic form:

(a)  $5^3 = 125$

(b)  $10^{-1} = 0.1$

(c)  $e^z = 4$

(d)  $6^0 = 1$

(e)  $2^{-3} = \frac{1}{8}$

(f)  $c^w = v$

7. Express these in exponential form:

(a)  $\log_5 625 = 4$

(b)  $\log_{10} 0.001 = -3$

(c)  $\ln e^2 = 2$

(d)  $\log_3 \frac{1}{81} = -4$

(e)  $\log_c v = w$

(f)  $\ln e^7 = 7$

8. Evaluate these logarithms without a calculator:

(a)  $\log_5 25$

(b)  $\log_4 64$

(c)  $\log_{10} 100000$

(d)  $\log_6 \frac{1}{216}$

(e)  $\log_{27} 9$

(f)  $\log_{25} 5$

9. Use the laws of logarithms to simplify:

(a)  $\log_c 4 + \log_c 9$

(b)  $\log_c 35 - \log_c 7$

(c)  $5 \log_c 2$

(d)  $\log_c m + 4 \log_c n - \log_c p$

(e)  $\frac{1}{4} \log_c 81 + \log_c 7$

(f)  $\log_c(z^2 - 25) - \log_c(z - 5)$  where  $z > 5$

10. Sketch the graphs of these logarithmic functions:

- (a)  $y = \log_5 x$
- (b)  $y = \ln x$
- (c)  $y = \log_5 x + 2$
- (d)  $y = \log_5(x - 2)$
- (e)  $y = -\log_5 x$
- (f)  $y = \log_5(-x)$  for  $x < 0$

## Section C: Solving Logarithmic Equations

11. Solve these logarithmic equations:

- (a)  $\log_5 x = 2$
- (b)  $\log_4(x + 2) = 3$
- (c)  $\log_{10}(4x - 3) = 1$
- (d)  $\ln(x - 4) = 0$
- (e)  $\log_3(x^2) = 4$
- (f)  $4\log_5 x = 8$

12. Solve these equations involving multiple logarithms:

- (a)  $\log_5 x + \log_5 4 = 3$
- (b)  $\log_c 15 - \log_c 5 = \log_c x$
- (c)  $\log_4 x + \log_4(x - 5) = 2$
- (d)  $\log_{10} x - \log_{10}(x - 5) = \log_{10} 4$
- (e)  $4\log_3 x = \log_3 27$
- (f)  $\log_2(x + 3) + \log_2(x - 3) = 4$

13. Solve these equations where the base is unknown:

- (a)  $\log_a 36 = 2$
- (b)  $\log_a \frac{1}{64} = -3$
- (c)  $\log_a 128 = \frac{7}{3}$
- (d)  $\log_a 625 = \frac{4}{3}$

14. Solve these quadratic logarithmic equations:

- (a)  $(\log_5 x)^2 = 4$
- (b)  $(\log_3 x)^2 - 5\log_3 x + 6 = 0$
- (c)  $\log^2 x - 5\log x + 6 = 0$  (base 10)
- (d)  $(\ln x)^2 - 7\ln x + 12 = 0$

15. Use the change of base formula to evaluate:

- (a)  $\log_4 9$  in terms of natural logarithms
- (b)  $\log_6 20$  in terms of common logarithms
- (c)  $\log_3 15$  using  $\ln$
- (d) Express  $\log_m n \times \log_n p \times \log_p m$

## Section D: Combined Exponential and Logarithmic Equations

16. Solve these mixed equations:

- (a)  $e^x = 9$
- (b)  $5^x = 20$
- (c)  $6 \times 2^x = 48$
- (d)  $3^{x-2} = 18$
- (e)  $e^{2x} - 5e^x + 6 = 0$
- (f)  $5^{2x} - 6 \times 5^x + 5 = 0$

17. Solve using substitution methods:

- (a)  $25^x - 5^{x+2} - 20 = 0$  (let  $y = 5^x$ )
- (b)  $8^x - 3 \times 2^x + 2 = 0$  (let  $u = 2^x$ )
- (c)  $e^{2x} - 8e^x + 15 = 0$  (let  $t = e^x$ )
- (d)  $\log^2 x - 5 \log x + 6 = 0$  (let  $z = \log x$ )

18. Find the exact solutions:

- (a)  $\ln x + \ln(x - 3) = \ln 10$
- (b)  $\log_2 x + \log_8 x = 4$
- (c)  $e^x + e^{-x} = 5$
- (d)  $4 \ln x = \ln(x + 12)$

19. Solve these equations involving both exponentials and logarithms:

- (a)  $x = \log_5(5^x + 4)$
- (b)  $e^{\ln x} = x + 5$
- (c)  $\ln(e^x - 3) = 3$
- (d)  $\log_4(4^x + 12) = x + 3$

20. Find the values of  $x$  for which:

- (a)  $5^x > 625$
- (b)  $\log_4 x < 2$
- (c)  $e^x \leq 20$
- (d)  $\ln x \geq 2$
- (e)  $\log_5(x - 3) > 1$
- (f)  $4^{x-2} < \frac{1}{16}$

## Section E: Exponential Growth and Decay

21. A population grows according to  $P = P_0 e^{kt}$  where  $P_0 = 1200$  and  $k = 0.05$  per year.

- (a) Find the population after 6 years
- (b) How long for the population to quadruple?
- (c) What is the percentage growth rate per year?
- (d) Find when the population reaches 4000
- (e) Calculate the population after 15 years

22. A radioactive substance decays according to  $A = A_0 e^{-\lambda t}$  where  $\lambda = 0.0462$  per year.
- (a) If initially there are 120g, find the amount after 12 years
  - (b) Calculate the half-life of the substance
  - (c) How long for 85% to decay?
  - (d) What percentage remains after 30 years?
  - (e) Find when only 15g remains
23. An investment grows at 7% compound interest per annum.
- (a) Write the growth formula
  - (b) How long to quadruple the investment?
  - (c) If £4000 is invested, find the value after 10 years
  - (d) How long for the investment to reach £20000?
  - (e) Compare with simple interest at 7% over 12 years
24. The temperature of a cooling object follows Newton's law:  $T = T_{\text{room}} + (T_0 - T_{\text{room}})e^{-kt}$
- (a) If room temperature is 18°C, initial temperature is 88°C, and  $k = 0.12$  per minute, find the temperature after 12 minutes
  - (b) How long for the object to cool to 35°C?
  - (c) Find the temperature after 30 minutes
  - (d) What happens as  $t \rightarrow \infty$ ?
  - (e) If the object cools to 55°C after 6 minutes, find  $k$
25. Carbon-14 dating uses the formula  $A = A_0 e^{-0.000121t}$  where  $t$  is in years.
- (a) Calculate the half-life of carbon-14
  - (b) If a sample has 35% of its original carbon-14, find its age
  - (c) How old is a sample with 8% remaining?
  - (d) What percentage remains after 20000 years?
  - (e) Find the age of a sample with ratio 0.65 of living organisms

## Section F: Logarithmic Modeling and Applications

26. The Richter scale for earthquake magnitude is given by  $M = \log_{10}(\frac{I}{I_0})$  where  $I$  is intensity.
- (a) If one earthquake has magnitude 8 and another has magnitude 5, compare their intensities
  - (b) An earthquake has intensity 50000 times the reference level. Find its magnitude
  - (c) How much more intense is a magnitude 7.5 earthquake than magnitude 5.5?
  - (d) Find the magnitude of an earthquake with intensity  $8 \times 10^5 I_0$
27. The pH scale is defined as  $\text{pH} = -\log_{10}[\text{H}^+]$  where  $[\text{H}^+]$  is hydrogen ion concentration.
- (a) Find the pH when  $[\text{H}^+] = 10^{-5}$  mol/L
  - (b) If  $\text{pH} = 1.5$ , find the hydrogen ion concentration
  - (c) Compare the acidity of solutions with pH 4 and pH 7
  - (d) Find the pH when  $[\text{H}^+] = 4.2 \times 10^{-3}$  mol/L
  - (e) If the concentration is halved, how does the pH change?
28. Sound intensity level in decibels is  $L = 10 \log_{10}(\frac{I}{I_0})$  where  $I_0 = 10^{-12}$  W/m<sup>2</sup>.

- (a) Find the decibel level when  $I = 10^{-4} \text{ W/m}^2$
  - (b) A sound has level 75 dB. Find its intensity
  - (c) How much more intense is 95 dB than 65 dB?
  - (d) Find the intensity of a 60 dB sound
  - (e) If intensity increases by factor 500, by how much do decibels increase?
29. The Michaelis-Menten equation in biochemistry is  $v = \frac{V_{\max}[S]}{K_m + [S]}$ .
- (a) Take logarithms to linearize when  $[S] \gg K_m$
  - (b) If  $V_{\max} = 120$ ,  $K_m = 6$ , find  $v$  when  $[S] = 18$
  - (c) Plot  $\log v$  against  $\log[S]$  for large  $[S]$
  - (d) Find  $[S]$  when  $v = \frac{3V_{\max}}{4}$
30. In information theory, entropy is  $H = -\sum p_i \log_2 p_i$ .
- (a) For a fair 8-sided die, calculate the entropy
  - (b) For a biased coin with  $P(H) = 0.8$ , find the entropy
  - (c) Find the entropy of a fair 16-sided die
  - (d) What probability distribution maximizes entropy for 5 outcomes?

## Section G: Advanced Functions and Transformations

31. Analyze the function  $f(x) = \ln(x - 4) + 1$ :
- (a) State the domain and range
  - (b) Find the x and y intercepts
  - (c) Identify any asymptotes
  - (d) Find  $f^{-1}(x)$
  - (e) Sketch both  $f(x)$  and  $f^{-1}(x)$
32. For the function  $g(x) = e^{4x-2} - 3$ :
- (a) Describe the transformations from  $y = e^x$
  - (b) State the domain and range
  - (c) Find the horizontal asymptote
  - (d) Solve  $g(x) = 0$
  - (e) Find  $g^{-1}(x)$
33. Consider  $h(x) = \log_4(16 - x^2)$ :
- (a) Find the domain of  $h(x)$
  - (b) Determine the range
  - (c) Find the maximum value and where it occurs
  - (d) Solve  $h(x) = 1$
  - (e) Sketch the graph of  $y = h(x)$
34. The function  $k(x) = me^{nx} + q$  passes through  $(0, 9)$ ,  $(1, 15)$ , and has horizontal asymptote  $y = 4$ .
- (a) Find the values of  $m$ ,  $n$ , and  $q$
  - (b) Write the equation of  $k(x)$

- (c) Find  $k(2)$
  - (d) Solve  $k(x) = 20$
  - (e) Find the domain and range of  $k(x)$
35. Investigate the function  $m(x) = \frac{\ln x}{x}$  for  $x > 0$ :
- (a) Find  $m'(x)$  and  $m''(x)$
  - (b) Locate any stationary points
  - (c) Determine the nature of stationary points
  - (d) Find the behavior as  $x \rightarrow 0^+$  and  $x \rightarrow \infty$
  - (e) Sketch the graph of  $y = m(x)$

## Section H: Systems and Complex Problems

36. Solve these simultaneous equations:

- (a) 
$$\begin{cases} y = 4^x \\ y = 8 - x \end{cases}$$
- (b) 
$$\begin{cases} \ln y = 4x \\ y = e^{x+3} \end{cases}$$
- (c) 
$$\begin{cases} \log_4 x + \log_4 y = 2 \\ x - y = 8 \end{cases}$$
- (d) 
$$\begin{cases} e^x + e^y = 10 \\ e^x - e^y = 6 \end{cases}$$

37. Find where these curves intersect:

- (a)  $y = e^x$  and  $y = \ln x$
- (b)  $y = 4^x$  and  $y = x^4$
- (c)  $y = \log x$  and  $y = 4 - x$
- (d)  $y = e^{-x}$  and  $y = x + 3$

38. Solve these differential equations:

- (a)  $\frac{dy}{dx} = ky$  where  $y(0) = y_0$
- (b)  $\frac{dP}{dt} = rP(1 - \frac{P}{K})$  (logistic growth)
- (c)  $\frac{dT}{dt} = -k(T - T_{\text{env}})$  (Newton's cooling)
- (d)  $\frac{dN}{dt} = -\lambda N$  (radioactive decay)

39. A bacteria culture follows logistic growth:  $P(t) = \frac{L}{1 + ae^{-kt}}$

- (a) If  $L = 1500$ ,  $P(0) = 60$ , and  $P(1) = 120$ , find  $a$  and  $k$
- (b) Find the population after 7 days
- (c) When does the population reach 750?
- (d) Find the maximum growth rate and when it occurs
- (e) Compare with exponential growth  $P = 60e^{rt}$

40. The Arrhenius equation in chemistry is  $k = Ae^{-E_a/(RT)}$  where  $k$  is reaction rate.

- (a) Take natural logarithms to linearize the equation

- (b) If at temperature 290K,  $k = 0.008$ , and at 340K,  $k = 0.12$ , find  $E_a/R$
- (c) Find the activation energy if  $R = 8.314 \text{ J/(mol}\cdot\text{K)}$
- (d) Predict the rate constant at 380K
- (e) At what temperature does the rate quadruple from 290K?

## Section I: Advanced Applications and Modeling

41. A pharmacokinetic model describes drug concentration:  $C(t) = \frac{D}{V}e^{-kt}$  where  $D$  is dose,  $V$  is volume of distribution,  $k$  is elimination rate.
  - (a) If  $D = 600 \text{ mg}$ ,  $V = 45 \text{ L}$ ,  $k = 0.12 \text{ h}^{-1}$ , find the initial concentration
  - (b) Calculate the concentration after 10 hours
  - (c) Find the half-life of the drug
  - (d) When does concentration drop to  $1.5 \text{ mg/L}$ ?
  - (e) Model repeated dosing every 6 hours
42. Economic growth follows  $Y(t) = Y_0e^{rt}$  where  $r$  is the growth rate.
  - (a) If GDP grows at 5% per year, how long to double?
  - (b) A country's GDP is £800 billion and grows to £1.4 trillion in 10 years. Find the growth rate
  - (c) Compare linear growth  $Y = Y_0(1 + rt)$  with exponential over 30 years
  - (d) Find when exponential growth overtakes linear with same initial rate
  - (e) Model with varying growth rate  $r(t) = r_0e^{-\gamma t}$
43. The spread of an epidemic follows  $I(t) = \frac{N}{1 + (N/I_0 - 1)e^{-rt}}$  (logistic model).
  - (a) If  $N = 12000$ ,  $I_0 = 15$ ,  $r = 0.18$  per day, find infections after 14 days
  - (b) When do infections peak?
  - (c) Find the maximum rate of spread
  - (d) Compare with exponential model  $I = I_0e^{rt}$  for early stages
  - (e) Model intervention reducing  $r$  by 60% after day 18
44. Weber-Fechner law relates stimulus and perception:  $P = k \log(S/S_0)$ .
  - (a) If quadrupling stimulus increases perception by 20 units, find  $k$
  - (b) Find perception when stimulus increases 12-fold
  - (c) A sound's loudness follows  $L = 10 \log_{10}(I/I_0)$ . Compare two sounds differing by 25 dB
  - (d) Model brightness perception where threshold  $S_0 = 0.08 \text{ lux}$
  - (e) Explain why geometric progressions produce arithmetic progressions in perception
45. Design an optimization problem involving exponentials:
  - (a) A company's profit is  $P(t) = 2000e^{0.06t} - 600t$  over  $t$  years
  - (b) Find when profit is maximized
  - (c) Calculate maximum profit
  - (d) Determine break-even points
  - (e) Model with discounting: present value =  $\frac{P(t)}{e^{rt}}$
  - (f) Find optimal time to sell considering 7% discount rate



**Answer Space**

Use this space for your working and answers.

**END OF TEST**

Total marks: 150

**For more resources and practice materials, visit:  
[stepupmaths.co.uk](http://stepupmaths.co.uk)**