

A Level Pure Mathematics

Practice Test 6: Integration

Instructions:

Answer all questions. Show your working clearly.
Calculators may be used unless stated otherwise.

Time allowed: 2 hours

Section A: Basic Integration - Polynomials

1. Find these indefinite integrals:

- (a) $\int (8x^2 + 3x - 9) dx$
- (b) $\int (6x^3 - 4x^2 + 7x + 5) dx$
- (c) $\int (7x^4 - 5x + 9) dx$
- (d) $\int (9x^2 - \frac{2}{7}x + 4) dx$
- (e) $\int (6x + 5)^2 dx$
- (f) $\int (5x - 3)(2x + 4) dx$

2. Integrate these functions involving negative and fractional powers:

- (a) $\int x^{-7} dx$
- (b) $\int (7x^{-1} + 4x^{\frac{5}{6}}) dx$
- (c) $\int \frac{6}{x^8} dx$
- (d) $\int \sqrt[3]{x} dx$
- (e) $\int \frac{7}{\sqrt{x}} dx$
- (f) $\int (5x^{\frac{8}{3}} - 6x^{-\frac{5}{6}}) dx$

3. Find these integrals by expanding first:

- (a) $\int \frac{6x^3 - 4x^2 + 5x}{x} dx$
- (b) $\int \frac{x^2 - 49}{x} dx$
- (c) $\int \frac{(4x - 3)^2}{x} dx$
- (d) $\int \frac{5x^3 + 64}{x^2} dx$

4. Evaluate these definite integrals:

- (a) $\int_2^4 (4x^2 + 3x - 2) dx$
- (b) $\int_1^6 (7x - 4) dx$
- (c) $\int_{-2}^3 x^3 dx$
- (d) $\int_9^{36} \sqrt{x} dx$

5. Find the function $f(x)$ given:

- (a) $f'(x) = 12x^2 + 9x - 7$ and $f(0) = 9$
- (b) $f'(x) = 16x - 7$ and $f(1) = 15$
- (c) $f''(x) = 14x - 12$, $f'(0) = 6$, and $f(0) = 8$
- (d) $f'(x) = \frac{6}{x^7}$ for $x > 0$ and $f(1) = 5$

Section B: Integration of Standard Functions

6. Integrate these exponential and logarithmic functions:

- (a) $\int 11e^x dx$
- (b) $\int 12e^x dx$
- (c) $\int e^{7x} dx$
- (d) $\int e^{-6x} dx$
- (e) $\int \frac{7}{x} dx$ for $x > 0$
- (f) $\int \frac{9}{x} dx$

7. Integrate these trigonometric functions:

- (a) $\int 11 \sin x dx$
- (b) $\int 10 \cos x dx$
- (c) $\int 12 \sin x dx$
- (d) $\int 7 \cos x dx$
- (e) $\int 7 \sec^2 x dx$
- (f) $\int 6 \operatorname{cosec}^2 x dx$

8. Find these integrals:

- (a) $\int (6 \sin x - 5 \cos x) dx$
- (b) $\int (7e^x + 4x^3) dx$
- (c) $\int (6e^x + 7 \cos x) dx$
- (d) $\int \left(\frac{6}{x} - 5x\right) dx$ for $x > 0$
- (e) $\int (7 \sin x + 5e^{-x}) dx$
- (f) $\int \left(6x^2 - \frac{7}{x^2}\right) dx$ for $x > 0$

9. Evaluate these definite integrals:

- (a) $\int_0^{6\pi} \cos x dx$
- (b) $\int_0^{\frac{\pi}{5}} \sin x dx$
- (c) $\int_0^6 e^x dx$
- (d) $\int_1^{e^6} \frac{1}{x} dx$
- (e) $\int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} \sec^2 x dx$
- (f) $\int_0^{\ln 7} e^{-x} dx$

10. Find the exact values:

- (a) $\int_0^{\frac{\pi}{6}} 7 \cos x dx$
- (b) $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \sin x dx$
- (c) $\int_0^{\ln 7} 5e^x dx$
- (d) $\int_1^{e^5} \frac{7}{x} dx$

Section C: Integration by Substitution

11. Use substitution to find these integrals:

- (a) $\int (7x - 4)^6 dx$
- (b) $\int (5x + 6)^5 dx$
- (c) $\int x(6x^2 - 5)^4 dx$
- (d) $\int x\sqrt{5x^2 + 7} dx$
- (e) $\int \frac{6x}{5x^2+4} dx$
- (f) $\int \frac{7x}{(5x^2-2)^2} dx$

12. Find these integrals using appropriate substitutions:

- (a) $\int \sin(7x - 4) dx$
- (b) $\int \cos(5x + \frac{\pi}{5}) dx$
- (c) $\int e^{7x-5} dx$
- (d) $\int e^{-7x} dx$
- (e) $\int \frac{1}{7x+4} dx$
- (f) $\int \frac{6}{5x-11} dx$

13. Use substitution for these more complex integrals:

- (a) $\int x^2(4x^3 - 9)^6 dx$
- (b) $\int \frac{x^2}{\sqrt{5x^3+4}} dx$
- (c) $\int xe^{6x^2} dx$
- (d) $\int \frac{\ln x}{6x} dx$
- (e) $\int \sin 6x \cos 5x dx$
- (f) $\int \csc 3x \cot 3x dx$

14. Evaluate these definite integrals using substitution:

- (a) $\int_0^4 x(3x^2 - 4)^4 dx$
- (b) $\int_0^{\frac{\pi}{10}} \sin 6x \cos 4x dx$
- (c) $\int_3^5 \frac{5x}{4x^2+3} dx$
- (d) $\int_0^4 xe^{5x^2} dx$

15. Find these integrals by recognizing the derivative pattern:

- (a) $\int \frac{12x-5}{6x^2-5x+3} dx$
- (b) $\int \frac{15x^2+10}{5x^3+10x-4} dx$
- (c) $\int \frac{6e^x}{e^x+5} dx$
- (d) $\int \frac{5\cos x}{\sin x} dx$

Section D: Integration by Parts

16. Use integration by parts to find:

- (a) $\int 6xe^x dx$
- (b) $\int 5x \sin x dx$
- (c) $\int 5x \cos x dx$
- (d) $\int x^2 e^{6x} dx$
- (e) $\int 6x \ln x dx$
- (f) $\int e^x \sin 5x dx$

17. Apply integration by parts to:

- (a) $\int 6 \ln x dx$
- (b) $\int x^6 \ln x dx$
- (c) $\int 5x \ln x dx$
- (d) $\int \ln(6x - 3) dx$
- (e) $\int 4x \sin^{-1} x dx$
- (f) $\int x^2 \sin 5x dx$

18. Find these integrals that may require multiple applications:

- (a) $\int x^2 e^{-6x} dx$
- (b) $\int x^2 \cos 5x dx$
- (c) $\int e^{6x} \cos 5x dx$
- (d) $\int e^{6x} \sin 5x dx$
- (e) $\int \cos(\ln 5x) dx$
- (f) $\int x^3 e^{6x} dx$

19. Evaluate these definite integrals:

- (a) $\int_0^6 xe^x dx$
- (b) $\int_0^{\frac{\pi}{5}} x \sin x dx$
- (c) $\int_1^{e^6} x \ln x dx$
- (d) $\int_0^{\frac{\pi}{10}} x \cos 5x dx$

20. Prove these reduction formulas using integration by parts:

- (a) $I_n = \int x^n e^{6x} dx = \frac{x^n e^{6x}}{6} - \frac{n}{6} I_{n-1}$
- (b) $I_n = \int \csc^n x dx = -\frac{\csc^{n-2} x \cot x}{n-1} + \frac{n-2}{n-1} I_{n-2}$ for $n \geq 2$
- (c) Use the first formula to find $\int x^6 e^{6x} dx$

Section E: Area Under Curves

21. Find the area under these curves:

- (a) $y = 6x^2$ from $x = 0$ to $x = 4$
- (b) $y = 7x - 3$ from $x = 1$ to $x = 4$
- (c) $y = x^3 - 5x$ from $x = 0$ to $x = 3$
- (d) $y = 5 \sin x$ from $x = 0$ to $x = \frac{\pi}{5}$

22. Calculate the area between the curve and the x-axis:

- (a) $y = x^2 - 36$ from $x = -6$ to $x = 6$
- (b) $y = x^3 - 25x$ from $x = -5$ to $x = 5$
- (c) $y = 5 \cos x$ from $x = 0$ to $x = 2\pi$
- (d) $y = e^x - 6$ from $x = 0$ to $x = \ln 7$

23. Find the area between these curves:

- (a) $y = 6x^2$ and $y = 24$ from $x = 0$ to $x = 2$
- (b) $y = x^2$ and $y = 6x - 5$ from $x = 1$ to $x = 5$
- (c) $y = \sin 5x$ and $y = \cos 4x$ from $x = 0$ to $x = \frac{\pi}{10}$
- (d) $y = 5e^x$ and $y = 10$ from $x = 0$ to $x = \ln 2$

24. Find the total area enclosed by:

- (a) $y = x^2 - 36$ and the x-axis
- (b) $y = x^3 - 49x$ and the x-axis
- (c) $y = 5 \sin x$ and $y = 0$ from $x = 0$ to $x = 2\pi$
- (d) $y = x^2 + 6x - 7$ and the x-axis

25. A region is bounded by $y = 6x^2$, $y = 0$, $x = 3$, and $x = 5$.

- (a) Calculate the area of the region
- (b) Find the x-coordinate of the centroid
- (c) Calculate the moment about the y-axis
- (d) Find the average value of $y = 6x^2$ over $[3, 5]$

Section F: Fundamental Theorem of Calculus

26. Use the fundamental theorem to evaluate:

- (a) $\frac{d}{dx} \int_0^x 6t^2 dt$
- (b) $\frac{d}{dx} \int_6^x \sin t dt$
- (c) $\frac{d}{dx} \int_0^{6x} e^t dt$
- (d) $\frac{d}{dx} \int_{5x}^{x^2} \cos t dt$

27. Find these derivatives:

- (a) $\frac{d}{dx} \int_0^x \sqrt{36 + t^2} dt$
- (b) $\frac{d}{dx} \int_x^7 \frac{6}{t} dt$
- (c) $\frac{d}{dx} \int_{\sin 5x}^{\cos 4x} t^5 dt$
- (d) $\frac{d}{dx} \int_0^{x^5} \sin(t^6) dt$

28. Given $M(x) = \int_5^x f(t) dt$ where f is continuous:

- (a) Prove that $M'(x) = f(x)$
- (b) If $f(x) = 6x^2 - 5$, find $M(x)$
- (c) Verify that $M'(x) = f(x)$ for your answer
- (d) Calculate $M(7) - M(6)$ and interpret geometrically

29. Solve these differential equations using antiderivatives:

- (a) $\frac{dy}{dx} = 12x^3 + 10x - 6$ with $y(0) = 7$
- (b) $\frac{dy}{dx} = 6e^x - \sin x$ with $y(0) = 5$
- (c) $\frac{d^2y}{dx^2} = 16x + 14$ with $y'(0) = 6$ and $y(0) = 5$
- (d) $\frac{dy}{dx} = \frac{6}{x}$ with $y(1) = 7$

30. For the function $m(x) = \int_6^x \frac{1}{t} dt$:

- (a) Find $m'(x)$
- (b) Show that $m(xy) = m(x) + m(y)$ for $x, y > 0$
- (c) Prove that $m(x^n) = n \cdot m(x)$ for $x > 0$ and integer n
- (d) Express $m(x)$ in terms of elementary functions

Section G: Volumes of Revolution

31. Find the volume when these curves are rotated about the x-axis:

- (a) $y = 6x$ from $x = 0$ to $x = 4$
- (b) $y = 5x^2$ from $x = 0$ to $x = 4$
- (c) $y = \sqrt{6x}$ from $x = 0$ to $x = 6$
- (d) $y = e^{6x}$ from $x = 0$ to $x = 1$

32. Calculate volumes of revolution about the x-axis:

- (a) $y = 5x - 1$ from $x = 0$ to $x = 4$
- (b) $y = x^2 - 5$ from $x = -3$ to $x = 3$
- (c) $y = 5 \sin x$ from $x = 0$ to $x = \frac{\pi}{4}$
- (d) $y = \frac{6}{x}$ from $x = 1$ to $x = 6$

33. Find volumes when rotated about the y-axis:

- (a) $x = 6y^2$ from $y = 0$ to $y = 3$
- (b) $x = \sqrt{6y}$ from $y = 0$ to $y = 6$
- (c) $x = e^{6y}$ from $y = 0$ to $y = 1$
- (d) $x = 6 \ln y$ from $y = 1$ to $y = e^6$

34. Use the washer method to find volumes:

- (a) Region between $y = 5x^2$ and $y = 20$ rotated about x-axis
- (b) Region between $y = 6x$ and $y = x^2$ rotated about x-axis
- (c) Region between $y = 5e^x$ and $y = 6$ from $x = 0$ to $x = \ln(\frac{6}{5})$ rotated about x-axis
- (d) Region between $y = \sqrt{6x}$ and $y = 5x$ rotated about y-axis

35. A solid has circular cross-sections. The radius at height h is $r(h) = \sqrt{49 - h^2}$ for $0 \leq h \leq 7$.

- (a) Set up the integral for the volume
- (b) Calculate the volume
- (c) Identify the shape of the solid
- (d) Find the surface area if this represents a hemisphere

Section H: Applications in Physics and Engineering

36. A particle moves with velocity $v(t) = 5t^2 - 12t + 7$ m/s.
- (a) Find the displacement from $t = 0$ to $t = 6$
 - (b) Calculate the total distance traveled
 - (c) Find the position function if $s(0) = 18$
 - (d) Determine when the particle changes direction
 - (e) Calculate the average velocity over $[0, 6]$
37. The acceleration of an object is $a(t) = 12t - 16$ m/s².
- (a) Find the velocity if $v(0) = 7$ m/s
 - (b) Find the position if $s(0) = 5$
 - (c) Calculate the displacement from $t = 2$ to $t = 4$
 - (d) Find when the object is at rest
 - (e) Determine the object's position at minimum velocity
38. Water flows into a tank at rate $R(t) = 15 - 3t$ liters per minute.
- (a) Find the total volume added in the first 4 minutes
 - (b) If the tank initially contains 40 liters, find $V(t)$
 - (c) Calculate the average flow rate over 4 minutes
 - (d) Find when the flow rate becomes zero
 - (e) Determine the maximum volume in the tank
39. The gravitational potential energy per unit mass is $\phi = -\frac{GM}{r}$ where G , M are constants.
- (a) Find gravitational field $g = -\frac{d\phi}{dr}$
 - (b) Calculate work done moving mass m from r_1 to r_2
 - (c) Find escape velocity using energy conservation
 - (d) Compare with surface gravity acceleration
40. The voltage across an inductor is $v(t) = L \frac{di}{dt}$ where L is inductance.
- (a) Find current $i(t)$ given $v(t) = V_0 e^{-t/\tau}$ and $i(0) = 0$
 - (b) Calculate energy stored $E = \int_0^\infty i^2 R dt$ in resistor
 - (c) Find time constant τ in terms of L and R
 - (d) Determine when current reaches 63% of final value

Section I: Advanced Applications and Techniques

41. The center of mass of a thin rod from $x = a$ to $x = b$ with density $\rho(x)$ is: $\bar{x} = \frac{\int_a^b x\rho(x) dx}{\int_a^b \rho(x) dx}$
- (a) Find the center of mass of a rod from $x = 0$ to $x = 7$ with density $\rho(x) = 6x + 5$
 - (b) Calculate the total mass of the rod
 - (c) Find the center of mass if density is $\rho(x) = e^{6x}$
 - (d) Compare with uniform density $\rho(x) = 6$
42. The moment of inertia about the x-axis is $I_x = \int y^2 dm$ where $dm = \rho dA$.

- (a) Find I_x for the region under $y = 6x^2$ from $x = 0$ to $x = 1$ with uniform density
 - (b) Calculate the radius of gyration $r_g = \sqrt{\frac{I_x}{M}}$
 - (c) Find the moment of inertia about the y-axis
 - (d) Analyze the distribution of mass relative to axes
43. Arc length of a curve $y = f(x)$ from $x = a$ to $x = b$ is: $L = \int_a^b \sqrt{1 + (f'(x))^2} dx$
- (a) Find the arc length of $y = 6x^2$ from $x = 0$ to $x = 1$
 - (b) Calculate the arc length of $y = \ln(6x)$ from $x = 1$ to $x = e$
 - (c) Find the perimeter of one arch of $y = 5 \sin x$
 - (d) Derive the formula using vector calculus
44. Surface area of revolution about x-axis is: $S = 2\pi \int_a^b y \sqrt{1 + (y')^2} dx$
- (a) Find the surface area when $y = 6x$ from $x = 0$ to $x = 4$ is rotated
 - (b) Calculate surface area for $y = \sqrt{6x}$ from $x = 0$ to $x = 6$
 - (c) Find the surface area of a cone with base radius $6R$ and height $5h$
 - (d) Use geometric formula to verify result
45. Economic applications of integration:
- (a) If marginal cost is $MC(x) = 7x + 13$, find total cost function given fixed costs of £350
 - (b) Calculate consumer surplus if demand is $p = 60 - 6x^2$ and price is £24
 - (c) Find producer surplus for supply curve $p = 5x^2 + 7$ at equilibrium price £22
 - (d) Determine optimal taxation to maximize government revenue
46. Probability density functions satisfy $\int_{-\infty}^{\infty} f(x) dx = 1$.
- (a) Find the constant h so that $f(x) = hx^7$ is a PDF on $[0, 1]$
 - (b) Calculate $P(0.3 \leq X \leq 0.8)$ for this distribution
 - (c) Find the first and third quartiles
 - (d) Calculate the interquartile range
47. Design an integration problem modeling thermodynamics:
- (a) Define a heat capacity function varying with temperature
 - (b) Set up integrals for enthalpy and entropy changes
 - (c) Calculate work done in thermodynamic processes
 - (d) Interpret results for engine efficiency
 - (e) Discuss applications in energy systems
48. Advanced computational integration:
- (a) Use the trapezoidal rule with $n = 14$ to approximate $\int_0^4 e^{-x^2/2} dx$
 - (b) Apply Simpson's rule with $n = 14$ to the same integral
 - (c) Compare with normal distribution tables
 - (d) Analyze convergence properties of methods
 - (e) Research Monte Carlo integration techniques

Answer Space

Use this space for your working and answers.

END OF TEST

Total marks: 150

**For more resources and practice materials, visit:
stepupmaths.co.uk**