

A Level Pure Mathematics

Practice Test 1: Vectors

Instructions:

Answer all questions. Show your working clearly.
Calculators may be used unless stated otherwise.

Time allowed: 2 hours

Section A: Vector Basics and Notation

1. Given vectors $\mathbf{a} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} -1 \\ 4 \\ 2 \end{pmatrix}$, calculate:
 - (a) $\mathbf{a} + \mathbf{b}$
 - (b) $\mathbf{a} - \mathbf{b}$
 - (c) $2\mathbf{a} + 3\mathbf{b}$
 - (d) $3\mathbf{a} - 2\mathbf{b}$
 - (e) $|\mathbf{a}|$ and $|\mathbf{b}|$
 - (f) A unit vector in the direction of \mathbf{a}
2. Express these vectors in component form:
 - (a) \overrightarrow{AB} where $A(2, 3, -1)$ and $B(5, 1, 4)$
 - (b) \overrightarrow{PQ} where $P(-1, 2, 3)$ and $Q(4, -2, 1)$
 - (c) The position vector of point C if $\overrightarrow{OC} = 3\mathbf{i} - 2\mathbf{j} + 5\mathbf{k}$
 - (d) \overrightarrow{BA} where $A(1, -2, 3)$ and $B(4, 1, -2)$
3. Given $\mathbf{p} = \mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$ and $\mathbf{q} = 2\mathbf{i} - \mathbf{j} + 4\mathbf{k}$:
 - (a) Find $|\mathbf{p}|$ and $|\mathbf{q}|$
 - (b) Calculate $\mathbf{p} + \mathbf{q}$ and $\mathbf{p} - \mathbf{q}$
 - (c) Find scalars α and β such that $\alpha\mathbf{p} + \beta\mathbf{q} = \begin{pmatrix} 0 \\ 5 \\ 2 \end{pmatrix}$
 - (d) Determine if \mathbf{p} and \mathbf{q} are parallel
4. Points A , B , and C have position vectors $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 4 \\ 1 \\ 2 \end{pmatrix}$, and $\mathbf{c} = \begin{pmatrix} 2 \\ 5 \\ 1 \end{pmatrix}$.
 - (a) Find vectors \overrightarrow{AB} and \overrightarrow{AC}
 - (b) Calculate the lengths $|AB|$ and $|AC|$

- (c) Find the position vector of the midpoint of BC
(d) Determine if triangle ABC is isosceles
5. Find the values of t for which these vectors are perpendicular:

(a) $\mathbf{u} = \begin{pmatrix} 2 \\ t \\ 1 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} t \\ 3 \\ -2 \end{pmatrix}$

(b) $\mathbf{p} = \begin{pmatrix} 1 \\ 2t \\ 3 \end{pmatrix}$ and $\mathbf{q} = \begin{pmatrix} 4 \\ -1 \\ t \end{pmatrix}$

(c) $\mathbf{r} = t\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and $\mathbf{s} = 3\mathbf{i} + t\mathbf{j} + 4\mathbf{k}$

Section B: Dot Product (Scalar Product)

6. Calculate the dot product of these vectors:

(a) $\mathbf{a} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 4 \\ 2 \\ -1 \end{pmatrix}$

(b) $\mathbf{p} = 3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and $\mathbf{q} = \mathbf{i} - 2\mathbf{j} + 4\mathbf{k}$

(c) $\mathbf{u} = \begin{pmatrix} 1 \\ 4 \\ -2 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix}$

(d) $\mathbf{r} = 2\mathbf{i} + 3\mathbf{j}$ and $\mathbf{s} = 4\mathbf{i} - \mathbf{j} + 2\mathbf{k}$

7. Find the angle between these pairs of vectors:

(a) $\mathbf{a} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$

(b) $\mathbf{p} = \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$ and $\mathbf{q} = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$

(c) $\mathbf{u} = 3\mathbf{i} + 4\mathbf{j}$ and $\mathbf{v} = \mathbf{i} + \mathbf{j} + \mathbf{k}$

(d) $\mathbf{r} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ and $\mathbf{s} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$

8. Use the dot product to verify these properties:

(a) $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$ (commutative)

(b) $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$ (distributive)

(c) $(k\mathbf{a}) \cdot \mathbf{b} = k(\mathbf{a} \cdot \mathbf{b})$ for scalar k

(d) $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$

9. Given vectors $\mathbf{a} = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$, and $\mathbf{c} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix}$:

(a) Show that \mathbf{a} and \mathbf{b} are perpendicular

(b) Find the component of \mathbf{c} in the direction of \mathbf{a}

(c) Calculate $|\mathbf{a} + \mathbf{b} + \mathbf{c}|$

- (d) Find the angle between $\mathbf{a} + \mathbf{b}$ and \mathbf{c}
10. A triangle has vertices at $A(1, 2, 3)$, $B(4, 1, 2)$, and $C(2, 3, 1)$.
- Find the vectors \overrightarrow{AB} and \overrightarrow{AC}
 - Calculate the angle $\angle BAC$
 - Find the area of triangle ABC
 - Determine if the triangle is right-angled

Section C: Cross Product (Vector Product)

11. Calculate the cross product of these vectors:

- $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}$
- $\mathbf{p} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}$ and $\mathbf{q} = \mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$
- $\mathbf{u} = \begin{pmatrix} 3 \\ 0 \\ -1 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix}$
- $\mathbf{r} = \mathbf{i} + \mathbf{j}$ and $\mathbf{s} = \mathbf{j} + \mathbf{k}$

12. Verify these properties of the cross product:

- $\mathbf{a} \times \mathbf{b} = -(\mathbf{b} \times \mathbf{a})$ (anti-commutative)
- $\mathbf{a} \times (\mathbf{b} + \mathbf{c}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}$ (distributive)
- $\mathbf{a} \times \mathbf{a} = \mathbf{0}$
- $|\mathbf{a} \times \mathbf{b}|^2 = |\mathbf{a}|^2|\mathbf{b}|^2 - (\mathbf{a} \cdot \mathbf{b})^2$

13. Find the area of the parallelogram spanned by:

- $\mathbf{a} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$
- $\mathbf{p} = 3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and $\mathbf{q} = \mathbf{i} - \mathbf{j} + 4\mathbf{k}$
- Vectors from origin to points $(2, 1, 3)$ and $(1, 4, 2)$
- \overrightarrow{AB} and \overrightarrow{AC} where $A(1, 0, 1)$, $B(2, 1, 3)$, $C(0, 2, 1)$

14. Given $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix}$:

- Calculate $\mathbf{a} \times \mathbf{b}$
- Verify that $\mathbf{a} \times \mathbf{b}$ is perpendicular to both \mathbf{a} and \mathbf{b}
- Find a unit vector perpendicular to both \mathbf{a} and \mathbf{b}
- Calculate the area of triangle with sides \mathbf{a} and \mathbf{b}

15. Use the scalar triple product $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ to find:

- The volume of parallelepiped with edges $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$, $\mathbf{c} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$
- Whether points $A(1, 2, 3)$, $B(2, 1, 1)$, $C(3, 2, 2)$, $D(1, 1, 1)$ are coplanar
- The volume of tetrahedron with vertices at $(0, 0, 0)$, $(1, 2, 1)$, $(2, 1, 3)$, $(1, 1, 2)$

Section D: Equations of Lines

16. Find the vector equation of the line:

- (a) Passing through $A(2, 1, 3)$ in direction $\begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$
- (b) Passing through points $P(1, 2, 0)$ and $Q(3, -1, 4)$
- (c) Through origin parallel to vector $2\mathbf{i} - 3\mathbf{j} + \mathbf{k}$
- (d) Through $(4, -1, 2)$ parallel to the line $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + t \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$

17. Convert these to parametric form:

- (a) $\mathbf{r} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} + t \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$
- (b) $\mathbf{r} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} + s \begin{pmatrix} 2 \\ -1 \\ 4 \end{pmatrix}$
- (c) Line through $(1, 2, 3)$ and $(4, 0, -1)$
- (d) $\mathbf{r} = (3 + 2t)\mathbf{i} + (1 - t)\mathbf{j} + (2 + 3t)\mathbf{k}$

18. Find where these lines intersect the coordinate planes:

- (a) $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + t \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$ and the xy -plane
- (b) $\mathbf{r} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} + s \begin{pmatrix} 1 \\ 2 \\ -3 \end{pmatrix}$ and the xz -plane
- (c) Line through $(2, 1, 4)$ and $(0, 3, 1)$ with the yz -plane

19. Determine if these pairs of lines intersect, are parallel, or are skew:

- (a) $L_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} + s \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$
- (b) $L_1 : \mathbf{r} = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} + t \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} + s \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix}$
- (c) Lines through $(1, 2, 3)$ to $(2, 4, 5)$ and $(0, 1, 1)$ to $(3, 2, 4)$

20. Find the shortest distance between:

- (a) Point $(2, 1, 3)$ and line $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$
- (b) Parallel lines $L_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + s \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$
- (c) Skew lines $L_1 : \mathbf{r} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + s \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$

Section E: Equations of Planes

21. Find the equation of the plane:

- (a) With normal vector $\begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ passing through $(1, 2, -1)$
- (b) Passing through points $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$
- (c) Containing the lines $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ and $\mathbf{r} = \begin{pmatrix} 2 \\ 2 \\ 0 \end{pmatrix} + s \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$
- (d) Parallel to vectors $\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$ through $(1, 1, 1)$

22. Convert between vector and Cartesian forms:

- (a) $\mathbf{r} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + s \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + t \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix}$ to Cartesian form
- (b) $2x - y + 3z = 6$ to vector form
- (c) $x + 2y - z = 4$ to parametric form
- (d) $\mathbf{r} \cdot \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} = 5$ to Cartesian form

23. Find where these planes intersect coordinate axes:

- (a) $3x + 2y - z = 6$
- (b) $x - 2y + 4z = 8$
- (c) $2x + y + 3z = 12$
- (d) $x + y + z = 3$

24. Determine the relationship between these planes:

- (a) $\Pi_1 : x + 2y - z = 3$ and $\Pi_2 : 2x + 4y - 2z = 6$
- (b) $\Pi_1 : 2x - y + 3z = 4$ and $\Pi_2 : x + y - z = 2$
- (c) $\Pi_1 : x + y + z = 1$ and $\Pi_2 : 2x + 2y + 2z = 3$
- (d) $\Pi_1 : x - y + 2z = 5$ and $\Pi_2 : 2x + y - z = 1$

25. Find the line of intersection of these planes:

- (a) $x + y + z = 3$ and $2x - y + z = 1$
- (b) $2x + y - 3z = 4$ and $x - 2y + z = 1$
- (c) $3x - y + 2z = 6$ and $x + 2y - z = 3$
- (d) $x + 2y + 3z = 6$ and $2x - y + z = 4$

Section F: Angles and Distances

26. Find the angle between these planes:

- (a) $x + 2y - z = 3$ and $2x - y + 2z = 5$
- (b) $3x + y - 2z = 6$ and $x - 3y + z = 4$

(c) $\mathbf{r} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 2$ and $\mathbf{r} \cdot \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} = 3$

(d) $2x + 3y + z = 7$ and $x - y + 4z = 2$

27. Calculate the distance from point to plane:

(a) Point $(2, 1, 3)$ to plane $x + 2y - 2z = 4$

(b) Point $(1, -1, 2)$ to plane $3x - y + 2z = 6$

(c) Point $(0, 0, 0)$ to plane $2x + 3y - z = 12$

(d) Point $(4, 1, -2)$ to plane $x - 3y + 2z = 5$

28. Find the angle between line and plane:

(a) Line $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ and plane $x + y + z = 6$

(b) Line through $(1, 0, 2)$ and $(3, 2, 1)$ with plane $2x - y + 3z = 4$

(c) Line $\mathbf{r} = \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix} + s \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ and plane $3x + y - z = 2$

29. Determine where these lines intersect planes:

(a) $\mathbf{r} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $x + y + 2z = 8$

(b) Line through $(1, 2, 3)$ and $(4, 0, 1)$ with plane $2x - y + z = 7$

(c) $\mathbf{r} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} + s \begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix}$ and $x - 2y + 3z = 10$

30. Find the reflection of point in plane:

(a) Point $(2, 1, 3)$ in plane $x + y - z = 1$

(b) Point $(1, -1, 2)$ in plane $2x - y + 2z = 6$

(c) Point $(0, 3, 1)$ in plane $x + 2y + z = 4$

Section G: Advanced Vector Geometry

31. A tetrahedron has vertices at $A(1, 0, 0)$, $B(0, 2, 0)$, $C(0, 0, 3)$, and $D(1, 1, 1)$.

(a) Find the volume of the tetrahedron

(b) Calculate the area of face ABC

(c) Find the equation of the plane containing face ABC

(d) Determine the perpendicular distance from D to plane ABC

(e) Verify the volume using the distance formula

32. Three forces $\mathbf{F}_1 = 3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$, $\mathbf{F}_2 = \mathbf{i} - 4\mathbf{j} + 3\mathbf{k}$, and $\mathbf{F}_3 = -2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$ act on a particle.

(a) Find the resultant force

(b) Calculate the magnitude of the resultant

(c) Find a fourth force needed for equilibrium

- (d) If the forces act at point $(1, 2, 1)$, find the moment about the origin
33. A regular tetrahedron has vertices at $(1, 1, 1)$, $(1, -1, -1)$, $(-1, 1, -1)$, and $(-1, -1, 1)$.
- (a) Verify that all edges have equal length
 - (b) Find the center of the tetrahedron
 - (c) Calculate the angle between any two faces
 - (d) Find the equation of the sphere circumscribing the tetrahedron
34. The position vectors of points P , Q , and R are \mathbf{p} , \mathbf{q} , and \mathbf{r} respectively.
- (a) Express the centroid G in terms of \mathbf{p} , \mathbf{q} , and \mathbf{r}
 - (b) Show that $\overrightarrow{PG} + \overrightarrow{QG} + \overrightarrow{RG} = \mathbf{0}$
 - (c) If S is the midpoint of QR , express \overrightarrow{PS} in terms of position vectors
 - (d) Prove that the medians of triangle PQR meet at the centroid
35. A line passes through point $A(2, 1, 3)$ and is perpendicular to the plane $2x - y + 3z = 6$.
- (a) Find the vector equation of the line
 - (b) Calculate where the line intersects the plane
 - (c) Find the foot of perpendicular from A to the plane
 - (d) Calculate the distance from A to the plane

Section H: Applications and Problem Solving

36. A parallelogram $ABCD$ has vertices $A(1, 2, 1)$, $B(3, 1, 4)$, and $C(5, 4, 2)$.
- (a) Find the coordinates of vertex D
 - (b) Calculate the area of the parallelogram
 - (c) Find the lengths of the diagonals
 - (d) Determine if the parallelogram is a rhombus
 - (e) Calculate the angle between the diagonals
37. An aircraft flies from airport $A(100, 200, 5)$ to airport $B(400, 150, 8)$ (coordinates in km).
- (a) Find the displacement vector \overrightarrow{AB}
 - (b) Calculate the distance traveled
 - (c) If the flight takes 2 hours, find the average velocity vector
 - (d) Find the bearing of B from A (projected onto horizontal plane)
 - (e) Calculate the angle of climb
38. A pyramid has square base with vertices at $(2, 2, 0)$, $(-2, 2, 0)$, $(-2, -2, 0)$, $(2, -2, 0)$ and apex at $(0, 0, 4)$.
- (a) Find the volume of the pyramid
 - (b) Calculate the area of each triangular face
 - (c) Find the total surface area
 - (d) Determine the angle between a triangular face and the base
 - (e) Find the equation of the plane containing one triangular face

39. Two particles move along lines $L_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} + s \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$.
- Show that the lines are skew
 - Find the shortest distance between the lines
 - If the particles start at $t = s = 0$ and move with constant speeds, when are they closest?
 - Calculate their closest approach distance
 - Find the common perpendicular to both lines
40. A sphere has center $C(2, -1, 3)$ and radius 5.
- Write the equation of the sphere
 - Find where the sphere intersects the plane $x + y + z = 4$
 - Determine if the line $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + t \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ intersects the sphere
 - Find the equation of the tangent plane at point $(7, -1, 3)$
 - Calculate the volume and surface area of the sphere

Section I: Advanced Topics and Modeling

41. Prove these vector identities:
- $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$ (vector triple product)
 - $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c})$
 - $|\mathbf{a} \times \mathbf{b}|^2 + |\mathbf{a} \cdot \mathbf{b}|^2 = |\mathbf{a}|^2 |\mathbf{b}|^2$ (Lagrange identity)
 - $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$
42. Three planes $\Pi_1 : x + 2y - z = 3$, $\Pi_2 : 2x - y + 3z = 5$, and $\Pi_3 : 3x + y + 2z = 8$ intersect.
- Find their common point of intersection
 - Calculate the angles between each pair of planes
 - Find the line of intersection of Π_1 and Π_2
 - Determine the volume of the tetrahedron formed by the three planes and the origin
 - Verify the intersection point lies on all three planes
43. A coordinate system undergoes rotation. The new basis vectors are: $\mathbf{e}'_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$, $\mathbf{e}'_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}$, $\mathbf{e}'_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$
- Verify these form an orthonormal basis
 - Express vector $\begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix}$ in the new coordinate system
 - Find the rotation matrix for this transformation
 - Calculate the angle of rotation about the z-axis

44. A crystal has lattice vectors $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 1 \\ \sqrt{3} \\ 0 \end{pmatrix}$, $\mathbf{c} = \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix}$.
- (a) Calculate the volume of the unit cell
 - (b) Find the angles between the lattice vectors
 - (c) Determine the reciprocal lattice vectors
 - (d) Calculate the density if each unit cell contains 4 atoms of mass 10^{-23} g
 - (e) Find the distance between parallel planes with Miller indices $(1, 1, 0)$
45. Design a vector-based model for a real-world application:
- (a) Choose a scenario involving 3D geometry (robotics, computer graphics, engineering)
 - (b) Define your coordinate system and relevant vectors clearly
 - (c) Set up vector equations describing the system
 - (d) Solve a specific problem using vector methods
 - (e) Discuss advantages of vector methods for your application
 - (f) Consider limitations and potential extensions of your model
46. A satellite orbits Earth in an elliptical path. At time t , its position vector is: $\mathbf{r}(t) = a \cos(\omega t)\mathbf{i} + b \sin(\omega t)\mathbf{j} + c\mathbf{k}$
- (a) Find the velocity vector $\mathbf{v}(t) = \frac{d\mathbf{r}}{dt}$
 - (b) Calculate the acceleration vector $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt}$
 - (c) Show that acceleration is always directed toward the origin
 - (d) Find the speed as a function of time
 - (e) Determine when the satellite is closest to Earth
 - (f) Calculate the angular momentum $\mathbf{L} = \mathbf{r} \times m\mathbf{v}$
47. Vector calculus in 3D space involves del operator $\nabla = \frac{\partial}{\partial x}\mathbf{i} + \frac{\partial}{\partial y}\mathbf{j} + \frac{\partial}{\partial z}\mathbf{k}$.
- (a) For scalar field $\phi(x, y, z) = x^2 + y^2 + z^2$, find $\nabla\phi$
 - (b) For vector field $\mathbf{F} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, calculate $\nabla \cdot \mathbf{F}$ (divergence)
 - (c) Find $\nabla \times \mathbf{F}$ (curl) for the same vector field
 - (d) Verify that $\nabla \times (\nabla\phi) = \mathbf{0}$ for any scalar field ϕ
 - (e) Show that $\nabla \cdot (\nabla \times \mathbf{F}) = 0$ for any vector field \mathbf{F}
48. Integration and applications:
- (a) Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$ where $\mathbf{F} = y\mathbf{i} + x\mathbf{j}$ and C is the curve from $(0, 0)$ to $(1, 1)$ along $y = x^2$
 - (b) Calculate the work done by force $\mathbf{F} = (x+y)\mathbf{i} + (x-y)\mathbf{j} + z\mathbf{k}$ moving a particle from $(0, 0, 0)$ to $(1, 1, 1)$
 - (c) Find the flux of $\mathbf{F} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ through the surface of unit cube
 - (d) Apply Green's theorem to evaluate $\oint_C (x^2 + y^2)dx + 2xy dy$ around unit circle

Answer Space

Use this space for your working and answers.

END OF TEST

Total marks: 150

**For more resources and practice materials, visit:
stepupmaths.co.uk**