A Level Pure Mathematics Practice Test 2: Exponentials and Logarithms

Instructions:

Answer all questions. Show your working clearly. Calculators may be used unless stated otherwise.

Time allowed: 2 hours

Section A: Laws of Indices and Exponential Functions

1. Simplify these expressions using laws of indices:

(a)
$$3^4 \times 3^2 \times 3^{-3}$$

(b)
$$\frac{5^6 \times 5^{-3}}{5^2}$$

(c)
$$(2^3)^4 \times 2^{-7}$$

(d)
$$\frac{(3^4)^2 \times 3^{-2}}{3^5}$$

(e)
$$(2^3 \times 2^{-2})^4$$

(f)
$$\frac{9^{3x} \times 9^{x+2}}{9^{4x-1}}$$

2. Solve these exponential equations:

(a)
$$3^x = 81$$

(b)
$$2^{x-1} = 32$$

(c)
$$7^{2x+1} = 49$$

(d)
$$9^x = \frac{1}{27}$$

(e)
$$16^x = 4^{x+3}$$

(f)
$$3^{2x} = 9^{x-2}$$

3. Express in the form a^x where a is a rational number:

(a)
$$\left(\frac{1}{3}\right)^x \times 9^x$$

(b)
$$\frac{16^x}{4^{3x}}$$

(c)
$$(8)^{\frac{x}{3}} \times (\frac{1}{2})^x$$

(d)
$$\frac{36^x \times 6^{-2x}}{216^{\frac{x}{3}}}$$

4. Sketch the graphs of these exponential functions, showing key features:

(a)
$$y = 3^x$$

(b)
$$y = (\frac{1}{3})^x$$

(c)
$$y = 2^x - 1$$

(d)
$$y = 3^{x+1}$$

- (e) $y = -3^x$
- (f) $y = 3^{-x}$
- 5. For the function $f(x) = 2e^x$:
 - (a) State the domain and range
 - (b) Find the y-intercept
 - (c) Describe the behavior as $x \to \infty$ and $x \to -\infty$
 - (d) Find f'(x) and comment on the gradient
 - (e) Sketch the graph, showing the tangent at (0,2)

Section B: Logarithmic Functions and Properties

- 6. Express these in logarithmic form:
 - (a) $3^2 = 9$
 - (b) $10^{-3} = 0.001$
 - (c) $e^y = 5$
 - (d) $7^0 = 1$
 - (e) $4^{-1} = \frac{1}{4}$
 - (f) $b^z = w$
- 7. Express these in exponential form:
 - (a) $\log_3 81 = 4$
 - (b) $\log_{10} 0.01 = -2$
 - (c) $\ln e = 1$
 - (d) $\log_7 \frac{1}{49} = -2$
 - (e) $\log_b y = z$
 - (f) $\ln e^5 = 5$
- 8. Evaluate these logarithms without a calculator:
 - (a) $\log_3 9$
 - (b) $\log_2 32$
 - (c) $\log_{10} 10000$
 - (d) $\log_4 \frac{1}{64}$
 - (e) $\log_8 4$
 - (f) $\log_{16} 2$
- 9. Use the laws of logarithms to simplify:
 - (a) $\log_b 7 + \log_b 2$
 - (b) $\log_b 24 \log_b 6$
 - (c) $4\log_b 3$
 - (d) $\log_b p + 3\log_b q \log_b r$
 - (e) $\frac{1}{3}\log_b 27 + \log_b 5$
 - (f) $\log_b(y^2 16) \log_b(y 4)$ where y > 4
- 10. Sketch the graphs of these logarithmic functions:

- (a) $y = \log_3 x$
- (b) $y = \ln x$
- (c) $y = \log_3 x 1$
- (d) $y = \log_3(x+1)$
- (e) $y = -\log_3 x$
- (f) $y = \log_3(-x)$ for x < 0

Section C: Solving Logarithmic Equations

- 11. Solve these logarithmic equations:
 - (a) $\log_3 x = 4$
 - (b) $\log_2(x-1) = 3$
 - (c) $\log_{10}(3x+2) = 1$
 - (d) ln(x+3) = 0
 - (e) $\log_4(x^2) = 3$
 - $(f) 3\log_2 x = 6$
- 12. Solve these equations involving multiple logarithms:
 - (a) $\log_3 x + \log_3 2 = 3$
 - (b) $\log_b 12 \log_b 3 = \log_b x$
 - (c) $\log_2 x + \log_2(x-3) = 2$
 - (d) $\log_{10} x \log_{10} (x 4) = \log_{10} 3$
 - (e) $3\log_4 x = \log_4 16$
 - (f) $\log_3(x+2) + \log_3(x-2) = 2$
- 13. Solve these equations where the base is unknown:
 - (a) $\log_a 25 = 2$
 - (b) $\log_a \frac{1}{27} = -3$
 - (c) $\log_a 64 = \frac{3}{2}$
 - (d) $\log_a 243 = \frac{5}{2}$
- 14. Solve these quadratic logarithmic equations:
 - (a) $(\log_3 x)^2 = 9$
 - (b) $(\log_2 x)^2 4\log_2 x + 3 = 0$
 - (c) $\log^2 x 4\log x + 3 = 0$ (base 10)
 - (d) $(\ln x)^2 6 \ln x + 8 = 0$
- 15. Use the change of base formula to evaluate:
 - (a) $\log_2 5$ in terms of natural logarithms
 - (b) $\log_7 15$ in terms of common logarithms
 - (c) $\log_4 12$ using ln
 - (d) Express $\log_p q \times \log_q r \times \log_r p$

Section D: Combined Exponential and Logarithmic Equations

- 16. Solve these mixed equations:
 - (a) $e^x = 7$
 - (b) $3^x = 15$
 - (c) $4 \times 3^x = 36$
 - (d) $7^{x+1} = 28$
 - (e) $e^{2x} 4e^x + 3 = 0$
 - (f) $3^{2x} 4 \times 3^x + 3 = 0$
- 17. Solve using substitution methods:
 - (a) $9^x 3^{x+1} 6 = 0$ (let $y = 3^x$)
 - (b) $16^x 5 \times 4^x + 4 = 0$ (let $u = 4^x$)
 - (c) $e^{2x} 7e^x + 10 = 0$ (let $t = e^x$)
 - (d) $\log^2 x 4\log x + 3 = 0$ (let $z = \log x$)
- 18. Find the exact solutions:
 - (a) $\ln x + \ln(x+2) = \ln 8$
 - (b) $\log_3 x + \log_9 x = 2$
 - (c) $e^x + e^{-x} = 4$
 - (d) $3 \ln x = \ln(x+8)$
- 19. Solve these equations involving both exponentials and logarithms:
 - (a) $x = \log_3(3^x + 2)$
 - (b) $e^{\ln x} = x + 3$
 - (c) $\ln(e^x 2) = 2$
 - (d) $\log_2(2^x + 6) = x + 2$
- 20. Find the values of x for which:
 - (a) $3^x > 81$
 - (b) $\log_2 x < 3$
 - (c) $e^x \le 15$
 - (d) $\ln x \ge 1$
 - (e) $\log_3(x-2) > 2$
 - (f) $2^{x+1} < \frac{1}{8}$

Section E: Exponential Growth and Decay

- 21. A population grows according to $P = P_0 e^{kt}$ where $P_0 = 800$ and k = 0.04 per year.
 - (a) Find the population after 8 years
 - (b) How long for the population to triple?
 - (c) What is the percentage growth rate per year?
 - (d) Find when the population reaches 3000
 - (e) Calculate the population after 25 years

- 22. A radioactive substance decays according to $A = A_0 e^{-\lambda t}$ where $\lambda = 0.0347$ per year.
 - (a) If initially there are 80g, find the amount after 15 years
 - (b) Calculate the half-life of the substance
 - (c) How long for 95% to decay?
 - (d) What percentage remains after 40 years?
 - (e) Find when only 8g remains
- 23. An investment grows at 8% compound interest per annum.
 - (a) Write the growth formula
 - (b) How long to triple the investment?
 - (c) If £3000 is invested, find the value after 12 years
 - (d) How long for the investment to reach £15000?
 - (e) Compare with simple interest at 8% over 15 years
- 24. The temperature of a cooling object follows Newton's law: $T = T_{\text{room}} + (T_0 T_{\text{room}})e^{-kt}$
 - (a) If room temperature is 25°C, initial temperature is 95°C, and k=0.15 per minute, find the temperature after 8 minutes
 - (b) How long for the object to cool to 40°C?
 - (c) Find the temperature after 45 minutes
 - (d) What happens as $t \to \infty$?
 - (e) If the object cools to 60° C after 4 minutes, find k
- 25. Carbon-14 dating uses the formula $A = A_0 e^{-0.000121t}$ where t is in years.
 - (a) Calculate the half-life of carbon-14
 - (b) If a sample has 25% of its original carbon-14, find its age
 - (c) How old is a sample with 12% remaining?
 - (d) What percentage remains after 15000 years?
 - (e) Find the age of a sample with ratio 0.75 of living organisms

Section F: Logarithmic Modeling and Applications

- 26. The Richter scale for earthquake magnitude is given by $M = \log_{10}(\frac{I}{I_0})$ where I is intensity.
 - (a) If one earthquake has magnitude 7 and another has magnitude 5, compare their intensities
 - (b) An earthquake has intensity 10000 times the reference level. Find its magnitude
 - (c) How much more intense is a magnitude 9 earthquake than magnitude 6?
 - (d) Find the magnitude of an earthquake with intensity $3 \times 10^6 I_0$
- 27. The pH scale is defined as $pH = -\log_{10}[H^+]$ where $[H^+]$ is hydrogen ion concentration.
 - (a) Find the pH when $[H^+] = 10^{-4} \text{ mol/L}$
 - (b) If pH = 3, find the hydrogen ion concentration
 - (c) Compare the acidity of solutions with pH 2 and pH 6
 - (d) Find the pH when $[H^{+}] = 3.5 \times 10^{-5} \text{ mol/L}$
 - (e) If the concentration triples, how does the pH change?
- 28. Sound intensity level in decibels is $L = 10 \log_{10}(\frac{I}{I_0})$ where $I_0 = 10^{-12}$ W/m².

- (a) Find the decibel level when $I = 10^{-5} \text{ W/m}^2$
- (b) A sound has level 85 dB. Find its intensity
- (c) How much more intense is 100 dB than 80 dB?
- (d) Find the intensity of a 40 dB sound
- (e) If intensity increases by factor 1000, by how much do decibels increase?
- 29. The Michaelis-Menten equation in biochemistry is $v = \frac{V_{\text{max}}[S]}{K_m + |S|}$.
 - (a) Take logarithms to linearize when $[S] >> K_m$
 - (b) If $V_{\text{max}} = 80$, $K_m = 8$, find v when [S] = 12
 - (c) Plot $\log v$ against $\log[S]$ for large [S]
 - (d) Find [S] when $v = \frac{V_{\text{max}}}{A}$
- 30. In information theory, entropy is $H = -\sum p_i \log_2 p_i$.
 - (a) For a fair 4-sided die, calculate the entropy
 - (b) For a biased coin with P(H) = 0.6, find the entropy
 - (c) Find the entropy of a fair 8-sided die
 - (d) What probability distribution maximizes entropy for 3 outcomes?

Section G: Advanced Functions and Transformations

- 31. Analyze the function $f(x) = \ln(x+3) 2$:
 - (a) State the domain and range
 - (b) Find the x and y intercepts
 - (c) Identify any asymptotes
 - (d) Find $f^{-1}(x)$
 - (e) Sketch both f(x) and $f^{-1}(x)$
- 32. For the function $g(x) = e^{3x+1} 5$:
 - (a) Describe the transformations from $y = e^x$
 - (b) State the domain and range
 - (c) Find the horizontal asymptote
 - (d) Solve g(x) = 0
 - (e) Find $g^{-1}(x)$
- 33. Consider $h(x) = \log_3(9 x^2)$:
 - (a) Find the domain of h(x)
 - (b) Determine the range
 - (c) Find the maximum value and where it occurs
 - (d) Solve h(x) = 2
 - (e) Sketch the graph of y = h(x)
- 34. The function $k(x) = pe^{qx} + r$ passes through (0,7), (1,12), and has horizontal asymptote y = 3.
 - (a) Find the values of p, q, and r
 - (b) Write the equation of k(x)

- (c) Find k(2)
- (d) Solve k(x) = 15
- (e) Find the domain and range of k(x)
- 35. Investigate the function $m(x) = x^2 \ln x$ for x > 0:
 - (a) Find m'(x) and m''(x)
 - (b) Locate any stationary points
 - (c) Determine the nature of stationary points
 - (d) Find the behavior as $x \to 0^+$ and $x \to \infty$
 - (e) Sketch the graph of y = m(x)

Section H: Systems and Complex Problems

36. Solve these simultaneous equations:

(a)
$$\begin{cases} y = 3^x \\ y = 6 - x \end{cases}$$

(b)
$$\begin{cases} \ln y = 3x \\ y = e^{x+2} \end{cases}$$

(c)
$$\begin{cases} \log_3 x + \log_3 y = 3 \\ x + y = 12 \end{cases}$$

(d)
$$\begin{cases} e^x + e^y = 8 \\ e^x - e^y = 4 \end{cases}$$

- 37. Find where these curves intersect:
 - (a) $y = e^x$ and $y = \ln x$
 - (b) $y = 3^x \text{ and } y = x^3$
 - (c) $y = \log x$ and y = 3 x
 - (d) $y = e^{-x}$ and y = x + 2
- 38. Solve these differential equations:
 - (a) $\frac{dy}{dx} = ky$ where $y(0) = y_0$
 - (b) $\frac{dP}{dt} = rP(1 \frac{P}{K})$ (logistic growth)
 - (c) $\frac{dT}{dt} = -k(T T_{\text{env}})$ (Newton's cooling)
 - (d) $\frac{dN}{dt} = -\lambda N$ (radioactive decay)
- 39. A bacteria culture follows logistic growth: $P(t) = \frac{L}{1 + ae^{-kt}}$
 - (a) If L = 800, P(0) = 40, and P(1) = 80, find a and k
 - (b) Find the population after 6 days
 - (c) When does the population reach 400?
 - (d) Find the maximum growth rate and when it occurs
 - (e) Compare with exponential growth $P = 40e^{rt}$
- 40. The Arrhenius equation in chemistry is $k = Ae^{-E_a/(RT)}$ where k is reaction rate.
 - (a) Take natural logarithms to linearize the equation

- (b) If at temperature 280K, k = 0.005, and at 320K, k = 0.08, find E_a/R
- (c) Find the activation energy if $R = 8.314 \text{ J/(mol \cdot K)}$
- (d) Predict the rate constant at 360K
- (e) At what temperature does the rate triple from 280K?

Section I: Advanced Applications and Modeling

- 41. A pharmacokinetic model describes drug concentration: $C(t) = \frac{D}{V}e^{-kt}$ where D is dose, V is volume of distribution, k is elimination rate.
 - (a) If D = 750 mg, V = 50 L, k = 0.08 h⁻¹, find the initial concentration
 - (b) Calculate the concentration after 12 hours
 - (c) Find the half-life of the drug
 - (d) When does concentration drop to 2 mg/L?
 - (e) Model repeated dosing every 8 hours
- 42. Economic growth follows $Y(t) = Y_0 e^{rt}$ where r is the growth rate.
 - (a) If GDP grows at 4% per year, how long to double?
 - (b) A country's GDP is £1.5 trillion and grows to £2.2 trillion in 8 years. Find the growth rate
 - (c) Compare linear growth $Y = Y_0(1 + rt)$ with exponential over 25 years
 - (d) Find when exponential growth overtakes linear with same initial rate
 - (e) Model with varying growth rate $r(t) = r_0 e^{-\beta t}$
- 43. The spread of an epidemic follows $I(t) = \frac{N}{1 + (N/I_0 1)e^{-rt}}$ (logistic model).
 - (a) If N = 15000, $I_0 = 20$, r = 0.15 per day, find infections after 12 days
 - (b) When do infections peak?
 - (c) Find the maximum rate of spread
 - (d) Compare with exponential model $I = I_0 e^{rt}$ for early stages
 - (e) Model intervention reducing r by 40% after day 25
- 44. Weber-Fechner law relates stimulus and perception: $P = k \log(S/S_0)$.
 - (a) If tripling stimulus increases perception by 15 units, find k
 - (b) Find perception when stimulus increases 8-fold
 - (c) A sound's loudness follows $L = 10 \log_{10}(I/I_0)$. Compare two sounds differing by 30 dB
 - (d) Model brightness perception where threshold $S_0 = 0.05$ lux
 - (e) Explain why multiplicative changes produce additive perception changes
- 45. Design an optimization problem involving exponentials:
 - (a) A company's profit is $P(t) = 1500e^{0.08t} 400t$ over t years
 - (b) Find when profit is maximized
 - (c) Calculate maximum profit
 - (d) Determine break-even points
 - (e) Model with discounting: present value = $\frac{P(t)}{e^{rt}}$
 - (f) Find optimal time to sell considering 6% discount rate

Answer Space

Use this space for your working and answers.

END OF TEST

Total marks: 150

For more resources and practice materials, visit: stepupmaths.co.uk